

Papua Expeditions check-list birds

Best of Papua birding expedition

Below we present a simple species-level check-list of birds potentially to be encountered on our Best of Papua birding expedition, which covers the following locations in Papua or Indonesian New Guinea: (1) Biak Island; (2) Lake Habbema alpine plateau and Ibele Valley in the Snow Mountains; (3) northern lowlands near Jayapura; (4) Arfak montane forests; (5) Waigeo Island; and (6) Sorong lowlands.

We define Papua or Indonesian New Guinea as the region encompassed by the western half of the subcontinental island of New Guinea up to the border with Papua New Guinea at 141° longitude, plus the following groups of satellite islands: (1) the Raja Ampat archipelago (until recently generally referred to as the West Papuan Islands), including the main islands of Waigeo, Batanta, Salawati, Misool, Kofiau and Gebe; (2) the Geelvink Islands (also known as the Schouten Islands), including the main islands of Biak-Supiori, Numfor and Yapen; and (3) the Aru Islands. We further follow the delineation of the New Guinea or 'Papuan' region as proposed initially by Mayr (1941), and perpetuated by Rand and Gilliard (1967), and Beehler *et al.* (1986). Classification, taxonomy and nomenclature generally follow Sibley and Monroe (1990; 1993), with detailed annotation of present deviations in species-level taxonomy and vernacular naming envisaged for future updates of this check-list. We indicate these author's orders, families, subfamilies and/or tribes, but omitted their suborders, infraorders, parvorders and/or superfamilies for the sake of simplicity. For taxa belonging to superspecies or allospecies, who's treatment at species level is not widely accepted, we have encoded this by the use of scientific trinomials and square bracketing of the binomen.

We list 428 bird species potentially to be encountered on our Best of Papua birding expedition. These include 34 Papua endemics, 220 regional New Guinea endemics, 157 widespread residents (at least six of which also have visiting migratory populations not included again in the ensuing category), 16 regular migrants, and a single recently introduced yet poorly established species.

Struthioniformes > Casuariidae > Casuariini

Cassowaries (1 genus / 2 species)

| | | |
|--------------------|-----------------------------------|--------------------|
| Dwarf Cassowary | <i>Casuarius bennetti</i> | Resident |
| Northern Cassowary | <i>Casuarius unappendiculatus</i> | New Guinea endemic |

Craciformes > Megapodiidae

Megapodes (3 genera / 8 species)

| | | |
|--------------------------|---|--------------------|
| Wattled Brush-turkey | <i>Aepyodius arfakianus</i> | New Guinea endemic |
| Bruijn's Brush-turkey | <i>Aepyodius bruijnii</i> | Papua endemic |
| Red-billed Talegalla | <i>Talegalla cuvieri</i> | Papua endemic |
| Brown-collared Talegalla | <i>Talegalla jobiensis</i> | New Guinea endemic |
| Orange-footed Megapode | <i>Megapodius [freycinet] reinwardt</i> | Resident |
| Dusky Megapode | <i>Megapodius [freycinet] freycinet</i> | Resident |
| Biak Megapode | <i>Megapodius [freycinet] geelvinkianus</i> | Papua endemic |
| New Guinea Megapode | <i>Megapodius [freycinet] affinis</i> | New Guinea endemic |

Galliformes > Phasianidae

Quails (2 genera / 2 species)

| | | |
|---------------------|--------------------------------|--------------------|
| Blue-breasted Quail | <i>Coturnix chinensis</i> | Resident |
| Snow Mountain Quail | <i>Anurophasis monorthonyx</i> | New Guinea endemic |

Anseriformes > Dendrocygnidae

Whistling-Ducks (1 genus / 2 species)

| | | |
|--------------------------|--------------------------------------|----------|
| Spotted Whistling-Duck | <i>Dendrocygna guttata</i> | Resident |
| Wandering Whistling-Duck | <i>Dendrocygna [bicolor] arcuata</i> | Resident |

Anseriformes > Anatidae > Anatinae

Typical Ducks (2 genera / 4 species)

| | | |
|--------------------|--------------------------|--------------------|
| Radjah Shelduck | <i>Tadorna radjah</i> | Resident |
| Salvadori's Teal | <i>Anas waigiuiensis</i> | New Guinea endemic |
| Pacific Black Duck | <i>Anas superciliosa</i> | Resident |
| Grey Teal | <i>Anas gracilis</i> | Resident |

Bucerotiformes > Bucerotidae

Typical Hornbills (1 genus / 1 species)

| | | |
|------------------|------------------------|----------|
| Blyth's Hornbill | <i>Aceros plicatus</i> | Resident |
|------------------|------------------------|----------|

Coraciiformes > Coraciidae

Typical Rollers (1 genus / 1 species)

| | | |
|------------|------------------------------|------------------|
| Dollarbird | <i>Eurystomus orientalis</i> | Resident/Migrant |
|------------|------------------------------|------------------|

Coraciiformes > Alcedinidae

Alcedinid Kingfishers (2 genera / 3 species)

| | | |
|---------------------------|-----------------------|----------|
| Azure Kingfisher | <i>Alcedo azurea</i> | Resident |
| Little Kingfisher | <i>Alcedo pusilla</i> | Resident |
| Variable Dwarf Kingfisher | <i>Ceyx lepidus</i> | Resident |

Coraciiformes > Dacelonidae

Halcyonid Kingfishers (6 genera / 12 species)

| | | |
|----------------------------------|---------------------------------------|--------------------|
| Rufous-bellied Kookaburra | <i>Dacelo gaudichaud</i> | New Guinea endemic |
| Shovel-billed Kookaburra | <i>Clytoceyx rex</i> | New Guinea endemic |
| Blue-black Kingfisher | <i>Todiramphus nigrocyaneus</i> | New Guinea endemic |
| Collared Kingfisher | <i>Todiramphus chloris</i> | Resident |
| Beach Kingfisher | <i>Todiramphus saurophaga</i> | Resident |
| Sacred Kingfisher | <i>Todiramphus sanctus</i> | Resident |
| Hook-billed Kingfisher | <i>Melidora macrorrhina</i> | New Guinea endemic |
| Yellow-billed Kingfisher | <i>Syma torotoro</i> | Resident |
| Mountain Kingfisher | <i>Syma megarhyncha</i> | New Guinea endemic |
| Common Paradise-Kingfisher | <i>Tanysiptera galatea</i> | Resident |
| Biak Paradise-Kingfisher | <i>Tanysiptera [galatea] riedelii</i> | Papua endemic |
| Red-breasted Paradise-Kingfisher | <i>Tanysiptera nympha</i> | New Guinea endemic |

Coraciiformes > Meropidae

Bee-eaters (1 genus / 2 species)

| | | |
|-----------------------|---------------------------|----------|
| Blue-tailed Bee-eater | <i>Merops philippinus</i> | Resident |
| Rainbow Bee-eater | <i>Merops ornatus</i> | Migrant |

Cuculiformes > Cuculidae

Old World Cuckoos (7 genera / 11 species)

| | | |
|-------------------------------|--|--------------------|
| Brush Cuckoo | <i>Cacomantis variolosus</i> | Resident/Migrant |
| Chestnut-breasted Cuckoo | <i>Cacomantis castaneiventris</i> | Resident |
| Fan-tailed Cuckoo | <i>Cacomantis flabelliformis</i> | Resident |
| Long-billed Cuckoo | <i>Rhamphomantis megarhynchus</i> | New Guinea endemic |
| Little Bronze-Cuckoo | <i>Chrysococcyx minutillus</i> | Resident |
| Rufous-throated Bronze-Cuckoo | <i>Chrysococcyx ruficollis</i> | New Guinea endemic |
| White-eared Bronze-Cuckoo | <i>Chrysococcyx meyeri</i> | New Guinea endemic |
| White-crowned Koel | <i>Caliechthrus leucolophus</i> | New Guinea endemic |
| Dwarf Koel | <i>Microdynamis parva</i> | New Guinea endemic |
| Asian Koel | <i>Eudynamis [scolopacea] scolopacea</i> | Resident |
| Channel-billed Cuckoo | <i>Scythrops novaehollandiae</i> | Migrant |

Cuculiformes > Centropodidae

Coucals (1 genus / 3 species)

| | | |
|----------------------|--------------------------|--------------------|
| Greater Black Coucal | <i>Centropus menbeki</i> | New Guinea endemic |
|----------------------|--------------------------|--------------------|

| | | |
|---------------------|-----------------------------|--------------------|
| Lesser Black Coucal | <i>Centropus bernsteini</i> | New Guinea endemic |
| Biak Coucal | <i>Centropus chalybeus</i> | Papua endemic |

Psittaciformes > Psittacidae

Parrots (21 genera / 37 species)

| | | |
|-------------------------------|-------------------------------------|--------------------|
| Black Lory | <i>Chalcopsitta atra</i> | Papua endemic |
| Brown Lory | <i>Chalcopsitta duivenbodei</i> | New Guinea endemic |
| Violet-necked Lory | <i>Eos squamata</i> | Resident |
| Black-winged Lory | <i>Eos cyanogenia</i> | Papua endemic |
| Dusky Lory | <i>Pseudeos fuscata</i> | New Guinea endemic |
| Rainbow Lorikeet | <i>Trichoglossus haematodus</i> | Resident |
| Goldie's Lorikeet | <i>Psitteuteles goldiei</i> | New Guinea endemic |
| Black-capped Lory | <i>Lorius lory</i> | New Guinea endemic |
| Pygmy Lorikeet | <i>Charmosyna wilhelminae</i> | New Guinea endemic |
| Red-fronted Lorikeet | <i>Charmosyna rubronotata</i> | New Guinea endemic |
| Red-flanked Lorikeet | <i>Charmosyna placentis</i> | New Guinea endemic |
| Fairy Lorikeet | <i>Charmosyna pulchella</i> | New Guinea endemic |
| Josephine's Lorikeet | <i>Charmosyna josephinae</i> | New Guinea endemic |
| Papuan Lorikeet | <i>Charmosyna papou</i> | New Guinea endemic |
| Whiskered Lorikeet | <i>Oreopsittacus arfaki</i> | New Guinea endemic |
| Yellow-billed Lorikeet | <i>Neopsittacus musschenbroekii</i> | New Guinea endemic |
| Orange-billed Lorikeet | <i>Neopsittacus pullicauda</i> | New Guinea endemic |
| Palm Cockatoo | <i>Probosciger aterrimus</i> | Resident |
| Sulphur-crested Cockatoo | <i>Cacatua galerita</i> | Resident |
| Yellow-capped Pygmy-Parrot | <i>Micropsitta keiensis</i> | Resident |
| Geelvink Pygmy-Parrot | <i>Micropsitta geelvinkiana</i> | Papua endemic |
| Buff-faced Pygmy-Parrot | <i>Micropsitta pusio</i> | New Guinea endemic |
| Red-breasted Pygmy-Parrot | <i>Micropsitta bruijnii</i> | Resident |
| Double-eyed Fig-Parrot | <i>Cyclopsitta diophthalma</i> | Resident |
| Salvadori's Fig-Parrot | <i>Psittaculirostris salvadorii</i> | Papua endemic |
| Brehm's Tiger-Parrot | <i>Psittacella brehmii</i> | New Guinea endemic |
| Painted Tiger-Parrot | <i>Psittacella picta</i> | New Guinea endemic |
| Modest Tiger-Parrot | <i>Psittacella modesta</i> | New Guinea endemic |
| Madarasz's Tiger-Parrot | <i>Psittacella madaraszii</i> | New Guinea endemic |
| Red-cheeked Parrot | <i>Geoffroyus geoffroyi</i> | Resident |
| Blue-collared Parrot | <i>Geoffroyus simplex</i> | New Guinea endemic |
| Great-billed Parrot | <i>Tanygnathus megalorhynchus</i> | Resident |
| Eclectus Parrot | <i>Eclectus roratus</i> | Resident |
| Pesquet's Parrot | <i>Psittirichas fulgidus</i> | New Guinea endemic |
| Moluccan King-Parrot | <i>Alisterus amboinensis</i> | Resident |
| Papuan King-Parrot | <i>Alisterus chloropterus</i> | New Guinea endemic |
| Orange-fronted Hanging-Parrot | <i>Loriculus aurantiifrons</i> | New Guinea endemic |

Apodiformes > Apodidae > Apodinae > Collocalini

Swiftlets (2 genera / 4 species)

| | | |
|-------------------|--------------------------------|--------------------|
| Glossy Swiftlet | <i>Collocalia esculenta</i> | Resident |
| Mountain Swiftlet | <i>Aerodramus hirundinacea</i> | New Guinea endemic |
| Uniform Swiftlet | <i>Aerodramus vanikorensis</i> | Resident |
| Papuan Swiftlet | <i>Aerodramus papuensis</i> | New Guinea endemic |

Apodiformes > Apodidae > Apodinae > Chaeturini

Needletails (1 genus / 1 species)

| | | |
|-------------------|------------------------------|--------------------|
| Papuan Needletail | <i>Mearnsia novaeguineae</i> | New Guinea endemic |
|-------------------|------------------------------|--------------------|

Apodiformes > Hemiprocnidae

Treeswifts (1 genus / 1 species)

| | | |
|----------------------|----------------------------|----------|
| Moustached Treeswift | <i>Hemiprocne mystacea</i> | Resident |
|----------------------|----------------------------|----------|

Strigiformes > Tytonidae

Barn and Grass Owls (1 genus / 2 species)

| | | |
|-----------|-------------------------|----------|
| Sooty Owl | <i>Tyto tenebricosa</i> | Resident |
|-----------|-------------------------|----------|

| | | |
|---|----------------------------------|--------------------|
| Eastern Grass Owl | <i>Tyto longimembris</i> | Resident |
| Strigiformes > Strigidae | | |
| Typical Owls (2 genera / 3 species) | | |
| Beccari's Scops Owl | <i>Otus [magicus] beccarii</i> | Papua endemic |
| Rufous Owl | <i>Ninox rufa</i> | Resident |
| Jungle Hawk-Owl | <i>Ninox theomacha</i> | New Guinea endemic |
| Strigiformes > Aegothelidae | | |
| Owlet-Nightjars (1 genus / 5 species) | | |
| Large Owlet-Nightjar | <i>Aegotheles insignis</i> | New Guinea endemic |
| Barred Owlet-Nightjar | <i>Aegotheles bennettii</i> | New Guinea endemic |
| Wallace's Owlet-Nightjar | <i>Aegotheles wallacii</i> | New Guinea endemic |
| Archbold's Owlet-Nightjar | <i>Aegotheles archboldi</i> | New Guinea endemic |
| Mountain Owlet-Nightjar | <i>Aegotheles albertisi</i> | New Guinea endemic |
| Strigiformes > Podargidae | | |
| Australian Frogmouths (1 genus / 2 species) | | |
| Papuan Frogmouth | <i>Podargus papuensis</i> | Resident |
| Marbled Frogmouth | <i>Podargus ocellatus</i> | Resident |
| Strigiformes > Eurostopodidae | | |
| 'Eared'-Nightjars (1 genus / 2 species) | | |
| Papuan Nightjar | <i>Eurostopodus papuensis</i> | New Guinea endemic |
| Archbold's Nightjar | <i>Eurostopodus archboldi</i> | New Guinea endemic |
| Strigiformes > Caprimulgidae > Caprimulginae | | |
| Nightjars (1 genus / 1 species) | | |
| Large-tailed Nightjar | <i>Caprimulgus macrurus</i> | Resident |
| Columbiformes > Columbidae | | |
| Pigeons, Doves (13 genera / 34 species) | | |
| Metallic Pigeon | <i>Columba vitiensis</i> | Resident |
| Slender-billed Cuckoo-Dove | <i>Macropygia amboinensis</i> | Resident |
| Black-billed Cuckoo-Dove | <i>Macropygia nigrirostris</i> | Resident |
| Great Cuckoo-Dove | <i>Reinwardtoena reinwardtii</i> | Resident |
| Emerald Dove | <i>Chalcophaps indica</i> | Resident |
| Stephan's Dove | <i>Chalcophaps stephani</i> | Resident |
| New Guinea Bronzewing | <i>Henicophaps albifrons</i> | New Guinea endemic |
| Nicobar Pigeon | <i>Caloenas nicobarica</i> | Resident |
| Cinnamon Ground-Dove | <i>Gallicolumba rufigula</i> | New Guinea endemic |
| Bronze Ground-Dove | <i>Gallicolumba beccarii</i> | Resident |
| Thick-billed Ground-Pigeon | <i>Trugon terrestris</i> | New Guinea endemic |
| Pheasant Pigeon | <i>Otidiphaps nobilis</i> | New Guinea endemic |
| Wompoo Fruit-Dove | <i>Ptilinopus magnificus</i> | Resident |
| Pink-spotted Fruit-Dove | <i>Ptilinopus perlatus</i> | New Guinea endemic |
| Ornate Fruit-Dove | <i>Ptilinopus ornatus</i> | New Guinea endemic |
| Orange-fronted Fruit-Dove | <i>Ptilinopus aurantifrons</i> | New Guinea endemic |
| Superb Fruit-Dove | <i>Ptilinopus superbus</i> | Resident |
| Coroneted Fruit-Dove | <i>Ptilinopus coronulatus</i> | New Guinea endemic |
| Beautiful Fruit-Dove | <i>Ptilinopus pulchellus</i> | New Guinea endemic |
| White-bibbed Fruit-Dove | <i>Ptilinopus rivoli</i> | Resident |
| Yellow-bibbed Fruit-Dove | <i>Ptilinopus solomonensis</i> | Resident |
| Claret-breasted Fruit-Dove | <i>Ptilinopus viridis</i> | Resident |
| Orange-bellied Fruit-Dove | <i>Ptilinopus iozonus</i> | New Guinea endemic |
| Dwarf Fruit-Dove | <i>Ptilinopus naina</i> | New Guinea endemic |
| Spice Imperial-Pigeon | <i>Ducula myristicivora</i> | Papua endemic |
| Purple-tailed Imperial-Pigeon | <i>Ducula rufigaster</i> | New Guinea endemic |
| Shining Imperial-Pigeon | <i>Ducula chalconota</i> | New Guinea endemic |
| Pinon Imperial-Pigeon | <i>Ducula pinon</i> | New Guinea endemic |

| | | |
|--------------------------|---------------------------------|--------------------|
| Collared Imperial-Pigeon | <i>Ducula mullerii</i> | New Guinea endemic |
| Banded Imperial-Pigeon | <i>Ducula zoeae</i> | New Guinea endemic |
| Pied Imperial-Pigeon | <i>Ducula [bicolor] bicolor</i> | Resident |
| Papuan Mountain-Pigeon | <i>Gymnophaps albertisii</i> | Resident |
| Western Crowned-Pigeon | <i>Goura cristata</i> | Papua endemic |
| Victoria Crowned-Pigeon | <i>Goura victoria</i> | New Guinea endemic |

Gruiformes > Rallidae

Rails, Gallinules, Coots (7 genera / 10 species)

| | | |
|----------------------------|---|--------------------|
| Chestnut Forest-Rail | <i>Rallina rubra</i> | New Guinea endemic |
| White-striped Forest-Rail | <i>Rallina leucospila</i> | Papua endemic |
| Red-necked Crake | <i>Rallina tricolor</i> | Resident |
| Buff-banded Rail | <i>Gallirallus philippensis</i> | Resident |
| Rufous-tailed Bush-hen | <i>Amaurornis [oliaceus] moluccanus</i> | Resident |
| Spotless Crake | <i>Porzana tabuensis</i> | Resident |
| White-browed Crake | <i>Porzana cinerea</i> | Resident |
| New Guinea Flightless Rail | <i>Megacrex inepta</i> | New Guinea endemic |
| Purple Swamphen | <i>Porphyrio porphyrio</i> | Resident |
| Eurasian Coot | <i>Fulica atra</i> | Resident |

Ciconiiformes > Scolopacidae > Scolopacinae

Woodcocks, Snipes (2 genera / 3 species)

| | | |
|---------------------|--|--------------------|
| New Guinea Woodcock | <i>Scolopax [saturata] rosenbergii</i> | New Guinea endemic |
| Latham's Snipe | <i>Gallinago hardwickii</i> | Migrant |
| Swinhoe's Snipe | <i>Gallinago megala</i> | Migrant |

Ciconiiformes > Scolopacidae > Tringinae

Godwits, Curlews, Sandpipers, Phalaropes (3 genera / 4 species)

| | | |
|----------------------|---------------------------|---------|
| Whimbrel | <i>Numenius phaeopus</i> | Migrant |
| Common Sandpiper | <i>Tringa hypoleucos</i> | Migrant |
| Grey-tailed Tattler | <i>Tringa brevipes</i> | Migrant |
| Red-necked Phalarope | <i>Phalaropus lobatus</i> | Migrant |

Ciconiiformes > Charadriidae > Charadriinae

Plovers, Dotterels, Lapwings (1 genus / 1 species)

| | | |
|-----------------------|------------------------|---------|
| Pacific Golden Plover | <i>Pluvialis fulva</i> | Migrant |
|-----------------------|------------------------|---------|

Ciconiiformes > Laridae > Larinae > Sternini

Terns (2 genera / 7 species)

| | | |
|---------------------|---------------------------|----------|
| Great Crested-Tern | <i>Sterna bergii</i> | Resident |
| Lesser Crested-Tern | <i>Sterna bengalensis</i> | Resident |
| Black-naped Tern | <i>Sterna sumatrana</i> | Resident |
| Common Tern | <i>Sterna hirundo</i> | Resident |
| Bridled Tern | <i>Sterna anaethetus</i> | Resident |
| Brown Noddy | <i>Anous stolidus</i> | Resident |
| Black Noddy | <i>Anous minutus</i> | Resident |

Ciconiiformes > Accipitridae > Pandioninae

Osprey (monotypic family)

| | | |
|--------|--------------------------|----------|
| Osprey | <i>Pandion haliaetus</i> | Resident |
|--------|--------------------------|----------|

Ciconiiformes > Accipitridae > Accipitrinae

Hawks, Eagles (10 genera / 15 species)

| | | |
|--------------------------|--|--------------------|
| Pacific Baza | <i>Aviceda subcristata</i> | Resident |
| Long-tailed Buzzard | <i>Henicopernis longicauda</i> | New Guinea endemic |
| Whistling Kite | <i>Haliastur sphenurus</i> | Resident |
| Brahminy Kite | <i>Haliastur indus</i> | Resident |
| White-bellied Fish-Eagle | <i>Haliaeetus leucogaster</i> | Resident |
| Papuan Harrier | <i>Circus [spilonotus] spilothorax</i> | New Guinea endemic |

| | | |
|-----------------------|--|--------------------|
| Variable Goshawk | <i>Accipiter [novaehollandiae] hiogaster</i> | Resident |
| Black-mantled Goshawk | <i>Accipiter melanochlamys</i> | New Guinea endemic |
| Grey-headed Goshawk | <i>Accipiter poliocephalus</i> | New Guinea endemic |
| Collared Sparrowhawk | <i>Accipiter cirrhocephalus</i> | Resident |
| Meyer's Goshawk | <i>Accipiter meyerianus</i> | Resident |
| Doria's Goshawk | <i>Megatriorchis doriae</i> | New Guinea endemic |
| New Guinea Eagle | <i>Harpyopsis novaeguineae</i> | New Guinea endemic |
| Gurney's Eagle | <i>Aquila gurneyi</i> | Resident |
| Little Eagle | <i>Hieraetus morphnoides</i> | Resident |

Ciconiiformes > Falconidae

True Falcons (1 genus / 3 species)

| | | |
|--------------------|--------------------------|------------------|
| Brown Falcon | <i>Falco berigora</i> | Resident |
| Australian Kestrel | <i>Falco cenchroides</i> | Resident/Migrant |
| Oriental Hobby | <i>Falco severus</i> | Resident |

Ciconiiformes > Sulidae

Boobies, Gannets (1 genus / 1 species)

| | | |
|-------------|-------------------------|---------|
| Brown Booby | <i>Sula leucogaster</i> | Migrant |
|-------------|-------------------------|---------|

Ciconiiformes > Anhingidae

Darters (1 genus / 1 species)

| | | |
|-------------------|---|----------|
| Australian Darter | <i>Anhinga [melanogaster] novaehollandiae</i> | Resident |
|-------------------|---|----------|

Ciconiiformes > Phalacrocoracidae

Cormorants (1 genus / 2 species)

| | | |
|------------------------|-----------------------------------|----------|
| Little Pied Cormorant | <i>Phalacrocorax melanoleucos</i> | Resident |
| Little Black Cormorant | <i>Phalacrocorax sulcirostris</i> | Resident |

Ciconiiformes > Ardeidae

Hérons, Bitterns, Egrets (9 genera / 10 species)

| | | |
|--------------------|-------------------------------|--------------------|
| Little Egret | <i>Egretta garzetta</i> | Resident/Migrant |
| Pacific Reef-Egret | <i>Egretta sacra</i> | Resident |
| Great-billed Heron | <i>Ardea sumatrana</i> | Resident |
| Great Egret | <i>Casmerodius albus</i> | Resident |
| Intermediate Egret | <i>Mesophoyx intermedia</i> | Resident |
| Cattle Egret | <i>Bubulcus ibis</i> | Resident |
| Striated Heron | <i>Butorides striatus</i> | Resident |
| Rufous Night-Heron | <i>Nycticorax caledonicus</i> | Resident |
| Forest Bittern | <i>Zonerodius heliosylus</i> | New Guinea endemic |
| Black Bittern | <i>Ixobrychus flavicollis</i> | Resident |

Ciconiiformes > Threskiornithidae

Ibises, Spoonbills (1 genus / 1 species)

| | | |
|-----------------------|-----------------------------|----------|
| Australian White Ibis | <i>Threskiornis molucca</i> | Resident |
|-----------------------|-----------------------------|----------|

Ciconiiformes > Fregatidae

Frigatebirds (1 genus / 1 species)

| | | |
|--------------------|----------------------|----------|
| Lesser Frigatebird | <i>Fregata ariel</i> | Resident |
|--------------------|----------------------|----------|

Ciconiiformes > Procellariidae > Procellariinae

Petrels, Shearwaters (2 genera / 2 species)

| | | |
|-------------------------|-------------------------------|---------|
| Streaked Shearwater | <i>Calonectris leucomelas</i> | Migrant |
| Wedge-tailed Shearwater | <i>Puffinus pacificus</i> | Migrant |

Passeriformes > Pittidae

Pittas (1 genus / 2 species)

| | | |
|-------------------|----------------------------|----------|
| Hooded Pitta | <i>Pitta sordida</i> | Resident |
| Red-bellied Pitta | <i>Pitta erythrogaster</i> | Resident |

Passeriformes > Climacteridae

Australo-Papuan Treecreepers (1 genus / 1 species)

| | | |
|--------------------|---------------------------|--------------------|
| Papuan Treecreeper | <i>Cornobates placens</i> | New Guinea endemic |
|--------------------|---------------------------|--------------------|

Passeriformes > Ptilonorhynchidae

Bowerbirds (4 genera / 5 species)

| | | |
|-------------------------|---|--------------------|
| White-eared Catbird | <i>Ailuroedus buccoides</i> | New Guinea endemic |
| Spotted Catbird | <i>Ailuroedus melanotis</i> | Resident |
| Archbold's Bowerbird | <i>Archboldia [papuensis] papuensis</i> | Papua endemic |
| Vogelkop Bowerbird | <i>Amblyornis inornatus</i> | Papua endemic |
| Fawn-breasted Bowerbird | <i>Chlamydera cerviniiventris</i> | Resident |

Passeriformes > Maluridae > Malurini

Fairywrens (2 genera / 3 species)

| | | |
|----------------------------|-------------------------------|--------------------|
| Orange-crowned Fairywren | <i>Clytomyias insignis</i> | New Guinea endemic |
| White-shouldered Fairywren | <i>Malurus alboscapulatus</i> | New Guinea endemic |
| Emperor Fairywren | <i>Malurus cyanocephalus</i> | New Guinea endemic |

Passeriformes > Meliphagidae

Honeyeaters, Macgregor's 'Bird of Paradise' (16 genera / 38 species)

| | | |
|----------------------------|---|--------------------|
| Red-throated Myzomela | <i>Myzomela eques</i> | New Guinea endemic |
| Dusky Myzomela | <i>Myzomela obscura</i> | Resident |
| Black Myzomela | <i>Myzomela nigrita</i> | New Guinea endemic |
| Red-collared Myzomela | <i>Myzomela rosenbergii</i> | New Guinea endemic |
| Olive Straightbill | <i>Timeliopsis fulvigula</i> | New Guinea endemic |
| Tawny Straightbill | <i>Timeliopsis griseigula</i> | New Guinea endemic |
| Long-billed Honeyeater | <i>Melilestes mearghynchus</i> | New Guinea endemic |
| Green-backed Honeyeater | <i>Glycichaera fallax</i> | Resident |
| Olive Honeyeater | <i>Lichmera argentauris</i> | Resident |
| Forest Honeyeater | <i>Meliphaga montana</i> | New Guinea endemic |
| Hill-forest Honeyeater | <i>Meliphaga orientalis</i> | New Guinea endemic |
| Scrub Honeyeater | <i>Meliphaga albonotata</i> | New Guinea endemic |
| Puff-backed Honeyeater | <i>Meliphaga aruensis</i> | New Guinea endemic |
| Mimic Honeyeater | <i>Meliphaga analoga</i> | New Guinea endemic |
| Yellow-gaped Honeyeater | <i>Meliphaga flavirictus</i> | New Guinea endemic |
| Black-throated Honeyeater | <i>Lichenostomus subfrenatus</i> | New Guinea endemic |
| Varied Honeyeater | <i>Lichenostomus [virescens] versicolor</i> | New Guinea endemic |
| Tawny-breasted Honeyeater | <i>Xanthotis flaviventer</i> | Resident |
| Spotted Honeyeater | <i>Xanthotis polygramma</i> | New Guinea endemic |
| Orange-cheeked Honeyeater | <i>Oreornis chrysogenys</i> | Papua endemic |
| Plain Honeyeater | <i>Pycnopygius ixoides</i> | New Guinea endemic |
| Marbled Honeyeater | <i>Pycnopygius cinereus</i> | New Guinea endemic |
| Streak-headed Honeyeater | <i>Pycnopygius stictocephalus</i> | New Guinea endemic |
| Meyer's Friarbird | <i>Philemon meyeri</i> | New Guinea endemic |
| New Guinea Friarbird | <i>Philemon [buceroides] novaeguineae</i> | Resident |
| Olive-streaked Honeyeater | <i>Ptiloprora meekiana</i> | New Guinea endemic |
| Rufous-sided Honeyeater | <i>Ptiloprora erythropleura</i> | Papua endemic |
| Black-backed Honeyeater | <i>Ptiloprora perstriata</i> | New Guinea endemic |
| Sooty Melidectes | <i>Melidectes fuscus</i> | New Guinea endemic |
| Short-bearded Melidectes | <i>Melidectes nouhuysi</i> | New Guinea endemic |
| Cinnamon-browed Melidectes | <i>Melidectes ochromelas</i> | New Guinea endemic |
| Vogelkop Melidectes | <i>Melidectes leucostephes</i> | Papua endemic |
| Belford's Melidectes | <i>Melidectes belfordi</i> | New Guinea endemic |
| Ornate Melidectes | <i>Melidectes torquatus</i> | New Guinea endemic |
| Western Smoky Honeyeater | <i>Melipotes gymnops</i> | Papua endemic |
| Smoky Honeyeater | <i>Melipotes fumigatus</i> | New Guinea endemic |

| | | |
|---|---|--------------------------------|
| MacGregor's 'Bird of Paradise' Brown-backed Honeyeater | <i>Macgregoria pulchra</i> <i>Ramsayornis modestus</i> | New Guinea endemic Resident |
|---|---|--------------------------------|

Passeriformes > Pardalotidae > Acanthizinae > Sericornithini

Mouse-warblers, Scrubwrens (2 genera / 10 species)

| | | |
|------------------------|--|--------------------|
| Rusty Mouse-warbler | <i>Crateroscelis murina</i> | New Guinea endemic |
| Mountain Mouse-warbler | <i>Crateroscelis robusta</i> | New Guinea endemic |
| Beccari's Scrubwren | <i>Sericornis [beccarii] beccarii</i> | Resident |
| Perplexing Scrubwren | <i>Sericornis [beccarii] virgatus</i> | New Guinea endemic |
| Large Scrubwren | <i>Sericornis nouhuysi</i> | New Guinea endemic |
| Vogelkop Scrubwren | <i>Sericornis [rufescens] rufescens</i> | Papua endemic |
| Buff-faced Scrubwren | <i>Sericornis [rufescens] perspicillatus</i> | New Guinea endemic |
| Grey-green Scrubwren | <i>Sericornis arfakianus</i> | New Guinea endemic |
| Papuan Scrubwren | <i>Sericornis papuensis</i> | New Guinea endemic |
| Pale-billed Scrubwren | <i>Sericornis spilodera</i> | New Guinea endemic |

Passeriformes > Pardalotidae > Acanthizinae > Acanthizini

Thornbills, Gerygones (2 genera / 8 species)

| | | |
|-------------------------|---|--------------------|
| Papuan Thornbill | <i>Acanthiza murina</i> | New Guinea endemic |
| Mountain Gerygone | <i>Gerygone cinerea</i> | New Guinea endemic |
| Green-backed Gerygone | <i>Gerygone chloronotus</i> | Resident |
| Fairy Gerygone | <i>Gerygone palpebrosa</i> | Resident |
| Yellow-bellied Gerygone | <i>Gerygone chrysogaster</i> | New Guinea endemic |
| Large-billed Gerygone | <i>Gerygone [magnirostris] magnirostris</i> | Resident |
| Biak Gerygone | <i>Gerygone [magnirostris] hypoxantha</i> | Papua endemic |
| Brown-breasted Gerygone | <i>Gerygone ruficollis</i> | New Guinea endemic |

Passeriformes > Eopsaltriidae

Australo-Papuan Robins (9 genera / 16 species)

| | | |
|------------------------|-------------------------------------|--------------------|
| Greater Ground-robin | <i>Amalocichla sclateriana</i> | New Guinea endemic |
| Lesser Ground-robin | <i>Amalocichla incerta</i> | New Guinea endemic |
| Torrent Robin | <i>Monachella muelleriana</i> | Resident |
| Lemon-bellied Flyrobin | <i>Microeca flavigaster</i> | Resident |
| Olive Flyrobin | <i>Microeca flavovirescens</i> | New Guinea endemic |
| Canary Flyrobin | <i>Microeca papuana</i> | New Guinea endemic |
| Garnet Robin | <i>Eugerygone rubra</i> | New Guinea endemic |
| Alpine Robin | <i>Petroica bivittata</i> | New Guinea endemic |
| Black-sided Robin | <i>Poecilodryas hypoleuca</i> | New Guinea endemic |
| Black-throated Robin | <i>Poecilodryas albonotata</i> | New Guinea endemic |
| White-winged Robin | <i>Peneothello sigillatus</i> | New Guinea endemic |
| Smoky Robin | <i>Peneothello cryptoleucus</i> | Papua endemic |
| Blue-grey Robin | <i>Peneothello cyanus</i> | New Guinea endemic |
| Ashy Robin | <i>Heteromyias albispectularis</i> | Resident |
| Green-backed Robin | <i>Pachycephalopsis hattamensis</i> | Papua endemic |
| White-eyed Robin | <i>Pachycephalopsis poliosoma</i> | New Guinea endemic |

Passeriformes > Orthonychidae

Logrunners (1 genus / 1 species)

| | | |
|----------------------|---|--------------------|
| New Guinea Logrunner | <i>Orthonyx [temminckii] novaeguineae</i> | New Guinea endemic |
|----------------------|---|--------------------|

Passeriformes > Pomatostomidae

Australo-Papuan Babblers (1 genus / 1 species)

| | | |
|----------------|------------------------------|--------------------|
| Rufous Babbler | <i>Pomatostomus isidorei</i> | New Guinea endemic |
|----------------|------------------------------|--------------------|

Passeriformes > Corvidae > Cinclosomatinae

Quail-thrushes, Whipbirds, Jewel-Babblers (2 genera / 3 species)

| | | |
|-----------------------|--------------------------------|--------------------|
| Papuan Whipbird | <i>Androphobus viridis</i> | New Guinea endemic |
| Spotted Jewel-Babbler | <i>Ptilorrhoa leucosticta</i> | New Guinea endemic |
| Blue Jewel-Babbler | <i>Ptilorrhoa caerulescens</i> | New Guinea endemic |

Passeriformes > Incertae sedis

| | | |
|-------|-----------------------|--------------------|
| Ifrit | <i>Ifrita kowaldi</i> | New Guinea endemic |
|-------|-----------------------|--------------------|

Passeriformes > Corvidae > Pachycephalinae > Neosittini

Sittellas (1 genus / 2 species)

| | | |
|-----------------|----------------------------------|--------------------|
| Varied Sittella | <i>Daphoenositta chrysoptera</i> | Resident |
| Black Sittella | <i>Daphoenositta miranda</i> | New Guinea endemic |

Passeriformes > Corvidae > Pachycephalinae > Pachycephalini

Whistlers, Shrike-thrushes, Pitohuis, Wattled Ploughbill (7 genera / 15 species)

| | | |
|-----------------------|----------------------------------|--------------------|
| Mottled Whistler | <i>Rhagologus leucostigma</i> | New Guinea endemic |
| Goldenface | <i>Pachycare flavogrisea</i> | New Guinea endemic |
| Rufous-naped Whistler | <i>Aleadryas rufinucha</i> | New Guinea endemic |
| Island Whistler | <i>Pachycephala phaionotus</i> | Resident |
| Grey Whistler | <i>Pachycephala simplex</i> | Resident |
| Golden Whistler | <i>Pachycephala pectoralis</i> | Resident |
| Sclater's Whistler | <i>Pachycephala soror</i> | New Guinea endemic |
| Lorentz's Whistler | <i>Pachycephala lorentzi</i> | New Guinea endemic |
| Regent Whistler | <i>Pachycephala schlegelii</i> | New Guinea endemic |
| Sooty Shrike-thrush | <i>Colluricincla umbrina</i> | New Guinea endemic |
| Little Shrike-thrush | <i>Colluricincla megarhyncha</i> | Resident |
| Variable Pitohui | <i>Pitohui kirhocephalus</i> | New Guinea endemic |
| Rusty Pitohui | <i>Pitohui ferrugineus</i> | New Guinea endemic |
| Black Pitohui | <i>Pitohui nigrescens</i> | New Guinea endemic |
| Wattled Ploughbill | <i>Eulacestoma nigropectus</i> | New Guinea endemic |

Passeriformes > Corvidae > Corvinae > Corvini

Crows (1 genus / 3 species)

| | | |
|-------------------|-----------------------------|--------------------|
| Brown-headed Crow | <i>Corvus fuscicapillus</i> | New Guinea endemic |
| Grey Crow | <i>Corvus tristis</i> | New Guinea endemic |
| Torresian Crow | <i>Corvus orru</i> | Resident |

Passeriformes > Corvidae > Corvinae > Paradisaeini

Birds of Paradise, Melampitta (13 genera / 24 species)

| | | |
|---------------------------------|--------------------------------|--------------------|
| Lesser Melampitta | <i>Melampitta lugubris</i> | New Guinea endemic |
| Crested Bird of Paradise | <i>Cnemophilus macgregorii</i> | New Guinea endemic |
| Loria's Bird of Paradise | <i>Cnemophilus loriae</i> | New Guinea endemic |
| Glossy-mantled Manucode | <i>Manucodia atra</i> | New Guinea endemic |
| Crinkle-collared Manucode | <i>Manucodia chalybata</i> | New Guinea endemic |
| Jobi Manucode | <i>Manucodia jobiensis</i> | New Guinea endemic |
| Long-tailed Paradigalla | <i>Paradigalla carunculata</i> | Papua endemic |
| Short-tailed Paradigalla | <i>Paradigalla brevicauda</i> | New Guinea endemic |
| Black Sicklebill | <i>Epimachus fastuosus</i> | New Guinea endemic |
| Brown Sicklebill | <i>Epimachus meyeri</i> | New Guinea endemic |
| Black-billed Sicklebill | <i>Epimachus albertsi</i> | New Guinea endemic |
| Pale-billed Sicklebill | <i>Epimachus bruijnii</i> | New Guinea endemic |
| Superb Bird of Paradise | <i>Lophorina superba</i> | New Guinea endemic |
| Western Parotia | <i>Parotia sefilata</i> | Papua endemic |
| Magnificent Riflebird | <i>Ptiloris magnificus</i> | Resident |
| Magnificent Bird of Paradise | <i>Cicinnurus magnificus</i> | New Guinea endemic |
| Wilson's Bird of Paradise | <i>Cicinnurus respublica</i> | Papua endemic |
| King Bird of Paradise | <i>Cicinnurus regius</i> | New Guinea endemic |
| Arfak Astrapia | <i>Astrapia nigra</i> | Papua endemic |
| Splendid Astrapia | <i>Astrapia splendidissima</i> | New Guinea endemic |
| King-of-Saxony Bird of Paradise | <i>Pteridophora alberti</i> | New Guinea endemic |
| Twelve-wired Bird of Paradise | <i>Seleucidis melanoleuca</i> | New Guinea endemic |
| Red Bird of Paradise | <i>Paradisaea rubra</i> | Papua endemic |
| Lesser Bird of Paradise | <i>Paradisaea minor</i> | New Guinea endemic |

Passeriformes > Corvidae > Corvinae > Artamini

Butcherbirds, Australian Magpie, Woodswallows, Peltops (3 genera / 6 species)

| | | |
|-----------------------------|------------------------------|--------------------|
| Hooded Butcherbird | <i>Cracticus cassicus</i> | New Guinea endemic |
| Black Butcherbird | <i>Cracticus quoyi</i> | Resident |
| White-breasted Wood-swallow | <i>Artamus leucorhynchus</i> | Resident |
| Great Wood-swallow | <i>Artamus maximus</i> | New Guinea endemic |
| Lowland Peltops | <i>Peltops blainvillii</i> | New Guinea endemic |
| Mountain Peltops | <i>Peltops montanus</i> | New Guinea endemic |

Passeriformes > Corvidae > Corvinae > Oriolini

Orioles, Cuckoo-shrikes, Trillers (4 genera / 13 species)

| | | |
|-----------------------------|---------------------------------|--------------------|
| Brown Oriole | <i>Oriolus szalayi</i> | New Guinea endemic |
| Stout-billed Cuckoo-shrike | <i>Coracina caeruleo-grisea</i> | New Guinea endemic |
| Yellow-eyed Cuckoo-shrike | <i>Coracina lineata</i> | Resident |
| Boyer's Cuckoo-shrike | <i>Coracina boyeri</i> | New Guinea endemic |
| White-bellied Cuckoo-shrike | <i>Coracina papuensis</i> | Resident |
| Hooded Cuckoo-shrike | <i>Coracina longicauda</i> | New Guinea endemic |
| Slender-billed Cicadabird | <i>Coracina tenuirostris</i> | Resident/Migrant |
| Black-shouldered Cicadabird | <i>Coracina incerta</i> | New Guinea endemic |
| Grey-headed Cuckoo-shrike | <i>Coracina schisticeps</i> | New Guinea endemic |
| New Guinea Cuckoo-shrike | <i>Coracina melas</i> | New Guinea endemic |
| Black-bellied Cuckoo-shrike | <i>Coracina montana</i> | New Guinea endemic |
| Golden Cuckoo-shrike | <i>Campochaera sloetii</i> | New Guinea endemic |
| Black-browed Triller | <i>Lalage atrovirens</i> | New Guinea endemic |

Passeriformes > Corvidae > Dicrurinae > Rhipidurini

Fantails (1 genus / 9 species)

| | | |
|-------------------------------|--------------------------------|--------------------|
| Willie-wagtail | <i>Rhipidura leucophrys</i> | Resident |
| Northern Fantail | <i>Rhipidura rufiventris</i> | Resident |
| Sooty Thicket-Fantail | <i>Rhipidura threnothorax</i> | New Guinea endemic |
| White-bellied Thicket-Fantail | <i>Rhipidura leucothorax</i> | New Guinea endemic |
| Black Fantail | <i>Rhipidura atra</i> | New Guinea endemic |
| Friendly Fantail | <i>Rhipidura albolimbata</i> | New Guinea endemic |
| Dimorphic Fantail | <i>Rhipidura brachyrhyncha</i> | New Guinea endemic |
| Rufous-backed Fantail | <i>Rhipidura rufidorsa</i> | New Guinea endemic |
| Rufous Fantail | <i>Rhipidura rufifrons</i> | Resident/Migrant |

Passeriformes > Corvidae > Dicrurinae > Dicrurini

Drongos (1 genus / 1 species)

| | | |
|-----------------|----------------------------|----------|
| Spangled Drongo | <i>Dicrurus bracteatus</i> | Resident |
|-----------------|----------------------------|----------|

Passeriformes > Corvidae > Dicrurinae > Monarchini

Monarchs, Magpie-larks (5 genera / 13 species)

| | | |
|--------------------------|--|--------------------|
| Black Monarch | <i>Monarcha axillaris</i> | New Guinea endemic |
| Island Monarch | <i>Monarcha cinerascens</i> | Resident |
| Spot-winged Monarch | <i>Monarcha guttulus</i> | New Guinea endemic |
| Hooded Monarch | <i>Monarcha manadensis</i> | New Guinea endemic |
| Biak Monarch | <i>Monarcha brehmii</i> | Papua endemic |
| Golden Monarch | <i>Monarcha chrysomela</i> | Resident |
| Frisled Monarch | <i>Arses [telescopthalmus] telescopthalmus</i> | Resident |
| Rufous-collared Monarch | <i>Arses [telescopthalmus] insularis</i> | New Guinea endemic |
| Biak Flycatcher | <i>Myiagra atra</i> | Papua endemic |
| Shining Flycatcher | <i>Myiagra alecto</i> | Resident |
| Yellow-breasted Boatbill | <i>Machaerirhynchus flaviventer</i> | Resident |
| Black-breasted Boatbill | <i>Machaerirhynchus nigripectus</i> | New Guinea endemic |
| Torrent-lark | <i>Grallina bruijni</i> | New Guinea endemic |

Passeriformes > Muscicapidae > Turdinae

True Thrushes (2 genera / 2 species)

| | | |
|----------------|----------------------------------|----------|
| Bassian Thrush | <i>Zoothera [dauma] lunulata</i> | Resident |
| Island Thrush | <i>Turdus poliocephalus</i> | Resident |

Passeriformes > Muscicapidae > Muscicapinae > Saxicolini

Chats (1 genus / 1 species)

| | | |
|---------------|-------------------------|----------|
| Pied Bushchat | <i>Saxicola caprata</i> | Resident |
|---------------|-------------------------|----------|

Passeriformes > Sturnidae > Sturnini

Starlings, Mynas (2 genera / 6 species)

| | | |
|----------------------|----------------------------|--------------------|
| Singing Starling | <i>Aplonis cantoroides</i> | Resident |
| Moluccan Starling | <i>Aplonis mysolensis</i> | Resident |
| Metallic Starling | <i>Aplonis metallica</i> | Resident |
| Long-tailed Starling | <i>Aplonis magna</i> | Papua endemic |
| Golden Myna | <i>Mino anais</i> | New Guinea endemic |
| Yellow-faced Myna | <i>Mino dumontii</i> | Resident |

Passeriformes > Hirundinidae > Hirundininae

Swallows (1 genus / 2 species)

| | | |
|-----------------|-------------------------|----------|
| Barn Swallow | <i>Hirundo rustica</i> | Migrant |
| Pacific Swallow | <i>Hirundo tahitica</i> | Resident |

Passeriformes > Pycnonotidae

Bulbuls (1 genus / 1 species)

| | | |
|---------------------|------------------------------|------------|
| Sooty-headed Bulbul | <i>Pycnonotus aurigaster</i> | Introduced |
|---------------------|------------------------------|------------|

Passeriformes > Cisticolidae

African Warblers (1 genus / 1 species)

| | | |
|-------------------------|-------------------------|----------|
| Golden-headed Cisticola | <i>Cisticola exilis</i> | Resident |
|-------------------------|-------------------------|----------|

Passeriformes > Zosteropidae

White-eyes (1 genus / 4 species)

| | | |
|-------------------------|------------------------------------|--------------------|
| Lemon-bellied White-eye | <i>Zosterops chloris</i> | Resident |
| Black-fronted White-eye | <i>Zosterops [atrifrons] minor</i> | New Guinea endemic |
| Biak White-eye | <i>Zosterops mysorensis</i> | Papua endemic |
| Capped White-eye | <i>Zosterops fuscicapillus</i> | New Guinea endemic |

Passeriformes > Sylviidae > Acrocephalinae

Gray's Grasshopper-Warbler, Reed-Warblers, Leaf-Warblers (3 genera / 3 species)

| | | |
|----------------------------|---|----------|
| Gray's Grasshopper-Warbler | <i>Locustella fasciolata</i> | Migrant |
| Clamorous Reed-Warbler | <i>Acrocephalus [stentoreus] stentoreus</i> | Resident |
| Island Leaf-Warbler | <i>Phylloscopus [trivirgatus] poliocephalus</i> | Resident |

Passeriformes > Sylviidae > Megalurinae

Grass-Warblers (1 genus / 2 species)

| | | |
|------------------|--|--------------------|
| Tawny Grassbird | <i>Megalurus [timoriensis] timoriensis</i> | Resident |
| Papuan Grassbird | <i>Megalurus [timoriensis] papuensis</i> | New Guinea endemic |

Passeriformes > Nectariniidae > Nectariniinae > Dicaeini

Flowerpeckers (1 genus / 2 species)

| | | |
|----------------------------|--|--------------------|
| Olive-crowned Flowerpecker | <i>Dicaeum [pectorale] pectorale</i> | Papua endemic |
| Red-capped Flowerpecker | <i>Dicaeum [pectorale] geelvinkianum</i> | New Guinea endemic |

Passeriformes > *Nectariniidae* > *Nectariniinae* > *Nectariniini*

Sunbirds (2 genera / 2 species)

| | | |
|----------------------|---------------------------|----------|
| Black Sunbird | <i>Leptocoma sericea</i> | Resident |
| Olive-backed Sunbird | <i>Cinnyris jugularis</i> | Resident |

Passeriformes > *Melanocharitidae* > *Melanocharitini*

Berrypeckers (1 genus / 4 species)

| | | |
|----------------------------|------------------------------------|--------------------|
| Black Berrypecker | <i>Melanocharis nigra</i> | New Guinea endemic |
| Lemon-breasted Berrypecker | <i>Melanocharis longicauda</i> | New Guinea endemic |
| Fan-tailed Berrypecker | <i>Melanocharis versteri</i> | New Guinea endemic |
| Streaked Berrypecker | <i>Melanocharis striativentris</i> | New Guinea endemic |

Passeriformes > *Melanocharitidae* > *Toxorhamphini*

Longbills (2 genera / 4 species)

| | | |
|------------------------|----------------------------------|--------------------|
| Green-crowned Longbill | <i>Toxorhamphus novaeguineae</i> | New Guinea endemic |
| Grey-winged Longbill | <i>Toxorhamphus poliopterus</i> | New Guinea endemic |
| Plumed Longbill | <i>Toxorhamphus iliolophus</i> | New Guinea endemic |
| Pygmy Longbill | <i>Oedistoma pygmaeum</i> | New Guinea endemic |

Passeriformes > *Paramythiidae*

Painted Berrypeckers (2 genera / 2 species)

| | | |
|---------------------|---------------------------|--------------------|
| Tit Berrypecker | <i>Oreocharis arfaki</i> | New Guinea endemic |
| Crested Berrypecker | <i>Paramythia montium</i> | New Guinea endemic |

Passeriformes > *Passeridae* > *Passerinae*

Sparrows (1 genus / 1 species)

| | | |
|-----------------------|------------------------|----------|
| Eurasian Tree Sparrow | <i>Passer montanus</i> | Resident |
|-----------------------|------------------------|----------|

Passeriformes > *Passeridae* > *Motacillinae*

Wagtails, Pipits (2 genera / 3 species)

| | | |
|----------------|--------------------------|--------------------|
| Yellow Wagtail | <i>Motacilla flava</i> | Migrant |
| Grey Wagtail | <i>Motacilla cinerea</i> | Migrant |
| Alpine Pipit | <i>Anthus gutturalis</i> | New Guinea endemic |

Passeriformes > *Passeridae* > *Estrildinae* > *Estrildini*

Estrildine Finches (3 genera / 8 species)

| | | |
|------------------------|---|--------------------|
| Mountain Firetail | <i>Oreostruthus fuliginosus</i> | New Guinea endemic |
| Blue-faced Parrotfinch | <i>Erythrura trichroa</i> | Resident |
| Papuan Parrotfinch | <i>Erythrura papuana</i> | New Guinea endemic |
| Streak-headed Munia | <i>Lonchura [tristissima] tristissima</i> | New Guinea endemic |
| Grand Munia | <i>Lonchura grandis</i> | New Guinea endemic |
| Hooded Munia | <i>Lonchura spectabilis</i> | Resident |
| Black-breasted Munia | <i>Lonchura teerinki</i> | Papua endemic |
| Snow Mountain Munia | <i>Lonchura montana</i> | New Guinea endemic |

References

- Beehler, B., Pratt, T. K. and Zimmerman, D.A. (1986). *Birds of New Guinea*. Princeton University Press: New Jersey.
- Mayr, E. (1941). *List of New Guinea Birds*. Amer. Mus. Nat. Hist. New York.
- Rand, A. L. and Gilliard, E. T. (1967). *Handbook of New Guinea Birds*. Weidenfeld & Nicholson: London.
- Sibley, C. G. and Monroe B. L. Jr. (1990). *Distribution and Taxonomy of Birds of the World*. Yale University Press: New Haven.
- Sibley, C. G. and Monroe B. L. Jr. (1993). *A Supplement to Distribution and Taxonomy of Birds of the World*. Yale University Press: New Haven.